


United Republic of Tanzania

Epidemiological Fact Sheet

on HIV/AIDS
and sexually
transmitted
infections



2000 Update




World Health
Organization

2 – United Republic of Tanzania

Country Information

Population pyramid, 1999



UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the working group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the working group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

The working group and its partners have established a framework standardizing the collection of data deemed important for a thorough understanding of the current status and trends of the epidemic, as well as patterns of risk and vulnerability in the population. Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed-upon indicators was not available for many countries in 1999. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the working group would like to encourage all programme managers as well as national and international experts to communicate additional information to the working group whenever such information becomes available. The working group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Indicators	Year	Estimate	Source
Total Population (thousands)	1999	32,793	UNPOP
Population Aged 15-49 (thousands)	1999	14,991	UNPOP
Annual Population Growth	1990-1998	2.9	UNPOP
% of Population Urbanized	1998	25	UNPOP
Average Annual Growth Rate of Urban Population	1990-1998	5.1	UNPOP
GNP Per Capita (US\$)	1997	210	World Bank
GNP Per Capita Average Annual Growth Rate	1996-1997	1.2	World Bank
Human Development Index Rank (HDI)	1999	156	UNDP
% Population Economic Active			
Unemployment Rate			
Total Adult Literacy Rate	1995	68	UNESCO
Adult Male Literacy Rate	1995	79	UNESCO
Adult Female Literacy Rate	1995	57	UNESCO
Male Secondary School Enrollment Ratio	1996	5.6	UNESCO
Female Secondary School Enrollment Ratio	1996	4.9	UNESCO
Crude Birth Rate (births per 1,000 pop.)	1999	41	UNPOP
Crude Death Rate (deaths per 1,000 pop.)	1999	15	UNPOP
Maternal Mortality Rate (per 100,000 live births)	1990	770	WHO
Life Expectancy at Birth	1998	48	UNPOP
Total Fertility Rate	1998	5.4	UNPOP
Infant Mortality Rate (per 1,000 live births)	1999	80	UNICEF/UNPOP

Contact address:
 UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance
 20, Avenue Appia
 CH-1211 Geneva 27
 Switzerland
 Fax: +41 22 791 4878
 email: surveillance@UNAIDS.org
<http://www.who.ch/emc/diseases/hiv>
<http://www.unaids.org>

Estimated number of people living with HIV/AIDS

In 1999 and during the first quarter of 2000, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1997 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalized epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 age range was used as the denominator in calculating adult HIV prevalence.

□ Estimated number of adults and children living with HIV/AIDS, end of 1999

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 1999

Adults and children	1300000		
Adults (15-49)	1200000	Adult rate (%)	8.09
Women (15-49)	670000		
Children (0-14)	59000		

□ Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 1999:

Deaths in 1999	140000
-----------------------	---------------

□ Estimated number of orphans

Estimated number of children who have lost their mother or both parents to AIDS (while they were under the age of 15) since the beginning of the epidemic:

Cumulative orphans	1100000
---------------------------	----------------

Estimated number of children who have lost their mother or both parents to AIDS and who were alive and under age 15 at the end of 1999:

Current living orphans	666697
-------------------------------	---------------

Assessment of epidemiological situation – United Republic of Tanzania

HIV information among antenatal clinic attendees has been available from Tanzania since the mid-1980s. In Dar es Salaam, the major urban area, HIV prevalence among antenatal clinic attendees tested increased from 4 percent in 1986 to 12 percent in 1995-96. Age detail is available for 1989, 1993 and 1995-96. In 1995-96, 7 percent of antenatal clinic attendees less than 20 years of age tested were HIV positive. Peak infection occurred among the 25-29 year old clinic attendees. Outside of Dar es Salaam, HIV information is available from sentinel surveillance reporting and from additional special studies. Median HIV prevalence increased from 3 percent in 1987 to 19 percent in 1998. In 1997, HIV prevalence among antenatal clinic attendees tested in the 14 reporting sites ranged from 4 to 44 percent.

Information on HIV prevalence among sex workers in Dar es Salaam has been available since the mid-1980s. HIV prevalence among sex workers tested increased from 29 percent in 1986 to 50 percent in 1993. Outside of Dar es Salaam, HIV information on sex workers is available from Kilimanjaro, Arusha, Moshi, Tanga, Dodoma, and Singida in 1988. Median HIV prevalence among sex workers tested in these areas was 19 percent, ranging from 8 to 32 percent. In 1992, 40 percent of sex workers tested in Mwanza town were HIV positive and in 1993, 61 percent of sex workers in Morogoro were HIV positive.

In Dar es Salaam, HIV prevalence among STD clinic patients tested increased from 13 percent in 1986 to 24 percent in 1994. In 1997, 13 percent of STD clinic patients tested were HIV positive. Outside of Dar es Salaam, most of the data on HIV prevalence among STD clinic patients comes from Mbeya. HIV prevalence among STD patients tested in Mbeya increased from 23 percent in 1988 to 27 percent in 1997. In Zanzibar, HIV prevalence increased from 5 percent in 1992 to 28 percent in 1993.

4 - United Republic of Tanzania

HIV sentinel surveillance

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV data base maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences is compiled. To provide for a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study on which the medians were calculated are printed at the end of this fact sheet.



The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and – where applicable – other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

□ HIV prevalence in selected populations in percent (for blood donors: 1/100 000)

Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Pregnant women	Major Urban Areas	N-sites			1		1	1	1	2	1	4	1	3				3
		Minimum			3.7		7.8	8.9	8.9	9.1	11	10.6	13.8	7.3				14.1
		Median			3.7		7.8	8.9	8.9	9.7	11	15.25	13.8	12.2				15.8
		Maximum			3.7		7.8	8.9	8.9	10.3	11	16.1	13.8	13.7				18.1
Pregnant women	Outside Major Urban Areas	N-sites				2	12	11	10	17	23	22	16	18	8	14	4	5
		Minimum				0.3	0.5	0.4	0.3	0	0.5	0.8	0.3	0	5.5	4	12.3	7.0
		Median				3.15	4.85	9.1	8.65	6.6	8	10.75	14	10.4	16.5	15.4	18.6	14.9
		Maximum				6	11	21.2	22.2	21	30.4	27.2	27.5	32.5	23.1	44.4	24	20.9
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Sex workers	Major Urban Areas	N-sites			1		1	1	1	1		1						
		Minimum			29		42.1	43.7	45	42.9		49.5						
		Median			29		42.1	43.7	45	42.9		49.5						
		Maximum			29		42.1	43.7	45	42.9		49.5						
Sex workers	Outside Major Urban Areas	N-sites					6				1	1						
		Minimum					7.5				39.8	60.9						
		Median					19.05				39.8	60.9						
		Maximum					31.9				39.8	60.9						
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Injecting drug users	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Injecting drug users	Outside Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI patients	Major Urban Areas	N-sites			4	2	2	2	1	5			1			1		
		Minimum			9.3	13.4	21.8	28.3	24.5	14.4			24			13.2		
		Median			12.7	20.75	24.8	34	24.5	18.5			24			13.2		
		Maximum			27.8	28.1	27.8	39.7	24.5	40.4			24			13.2		
STI patients	Outside Major Urban Areas	N-sites				1	2	1	1	1	2	1				9		
		Minimum				20	22.5	21.4	27.5	32	5.2	28.2				17		
		Median				20	23.05	21.4	27.5	32	19.8	28.2				27		
		Maximum				20	23.6	21.4	27.5	32	34.3	28.2				43		
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Blood Donors	National	N-sites																
		Minimum																
		Median																
		Maximum																
Blood Donors	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Men having sex with men	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																

Maps of HIV sentinel sites

Mapping the geographical distribution of HIV sentinel sites for different population groups may assist interpreting both the national coverage of the HIV surveillance system and explaining differences in levels and trends of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the UNICEF/WHO HealthMap Programme, has produced maps showing the location and HIV prevalence of HIV sentinel sites in relation to population density, major urban areas and communication routes. Maps illustrate separately the most recent results from HIV sentinel surveillance in pregnant women and in sub-populations at higher risk of HIV infection.



The boundaries and names shown and the designations used on these maps do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2000, all rights reserved.

6 - United Republic of Tanzania

Reported AIDS cases

AIDS cases by year of reporting

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	Total	Unkn
0	0	0	0	3	106	295	1121	2937	4839	5096	11106	18692	15871	13506	5873	4722	8426	10592	8867		112052	0

Date of last report: 11/Aug/99

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases is aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of HAART (Highly Active Anti-Retroviral Therapy).

AIDS cases by mode of transmission

Hetero: Heterosexual contacts.

Homo/Bi: Homosexual contacts between men.

IDU: Injecting drug use. This transmission category also includes cases in which other high-risk behaviours were reported, in addition to injection of drugs.

Blood: Blood and blood products.

Perinatal: Vertical transmission during pregnancy, birth or breastfeeding.

NS: Not specified/unknown.

Sex	Trans. Group	<96	1996	1997	1998	1999	Unkn	Total	%
All	Total								
	Hetero			9915					
	Homo/Bi								
	IDU								
	Blood			529					
	Perinatal			137					
	Other Known								
	Unknown								
Male	Total								
	Hetero			4769					
	Homo/Bi								
	IDU								
	Blood			267					
	Perinatal			70					
	Other Known								
	Unknown								
Female	Total								
	Hetero			4868					
	IDU								
	Blood			245					
	Perinatal			59					
	Other Known								
	Unknown								
NS	Total								
	Hetero			278					
	IDU								
	Blood			17					
	Perinatal			8					
	Other Known								
	Unknown								

Aids cases by age and sex

Sex	Age	<96	1996	1997	1998	1999	Unkn.	Total	%
All	All	82642	8426	10592	8867			110527	100.0
	0-4	3802	253	434	306			4795	4.3
	5-14	827	135	254	138			1354	1.2
	15-19	3141	169	223	194			3727	3.4
	20-29	31487	2300	3379	2367			39533	35.8
	30-39	28016	2485	4014	2816			37331	33.8
	40-49	9669	1019	1727	1157			13572	12.3
	50-59	2480	236	434	335			3485	3.2
	60+	661	59	106	77			903	0.8
	NS	2480	1778	21	1477			5756	5.2
Male	All	41404	3033	4999	3628			53064	100.0
	0-4	2070	121	215	149			2555	4.8
	5-14	373	61	115	69			618	1.2
	15-19	663	33	25	54			775	1.5
	20-29	11883	722	1130	850			14585	27.5
	30-39	15941	1234	2000	1485			20660	38.9
	40-49	6790	634	1130	736			9290	17.5
	50-59	1905	167	295	213			2580	4.9
	60+	538	40	70	58			706	1.3
	NS	1242	21	15	14			1292	2.4
Female	All	41238	3539	5307	3657			53741	100.0
	0-4	1773	117	202	143			2235	4.2
	5-14	495	71	112	61			739	1.4
	15-19	2474	127	191	132			2924	5.4
	20-29	19547	1557	2186	1486			24776	46.1
	30-39	12041	1196	1921	1289			16447	30.6
	40-49	2887	364	536	399			4186	7.8
	50-59	619	64	122	108			913	1.7
	60+	165	11	32	19			227	0.4
	NS	1237	32	5	20			1294	2.4
NS	All	0	1854	286	1582			3722	100.0
	0-4	0	7	18	14			39	1.0
	5-14	0	6	31	8			45	1.2
	15-19	0	6	7	8			21	0.6
	20-29	0	24	63	31			118	3.2
	30-39	0	43	101	42			186	5.0
	40-49	0	9	54	22			85	2.3
	50-59	0	4	9	14			27	0.7
	60+	0	6	0	0			6	0.2
	NS	0	1750	5	1443			3198	85.9

Curable Sexually Transmitted Infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STI are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STI facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Significant also is the observation of a sharp decline in the concentration of HIV in the genital secretions when the infection is treated. Monitoring trends in STI can provide valuable information on the sexual transmission of HIV as well as the impact of behavioural interventions, such as promotion of condom use.

Clinical services offering STI care are an important access point for people at high risk for both AIDS and STI, not only for diagnosis and treatment but also for information and education. Therefore, control and prevention of STI have been recognized as a major strategy in the prevention of HIV infection and ultimately AIDS. One of the cornerstones of STI control is adequate management of patients with symptomatic STIs. This includes diagnosis, treatment and individual health education and counselling on disease prevention and partner notification. Consequently, monitoring different components of STI control can also provide information on HIV prevention within a country.

☐ Estimated incidence and prevalence of curable STIs

STI's	Year	Incidence			Year	Prevalence		
		Male	Female	All		Male	Female	All
Chlamydia trach.								
Gonorrhoea								
Syphilis								
Trichomonas								

Comments:

Source:

☐ STI Incidence, men

Prevention Indicator 9: Proportion of men aged 15-49 years who reported episodes of urethritis in the last 12 months.

Year	Area	Age	Rate	N=
1996	All	15-49	2.2	

Comments:

Sources: Tanzania Demographic Health Surveys TDHS, 1996

☐ STI Prevalence, women

Prevention Indicator 8: Proportion of pregnant women aged 15-24 years attending antenatal clinics whose blood has been screened with positive serology for syphilis.

Year	Area	Age	Rate	N=
1996	All	15-24	5.3	1615

Comments:

Sources: National AIDS Control programme HIV/AIDS/STD Surveillance report no. 11

☐ STI Case management (counselled)

Prevention Indicator 7: Proportion of people presenting with STD or for STD care in health facilities who received basic advice on condoms and on partner notification.

Year	Area	Age	Rate	N=
1994	All		82.2	

Comments:

Sources: PPI 6&7 Survey. National AIDS Control Programme/WHO 1994

☐ STI Case management (treatments)

Prevention Indicator 6: Proportion of people presenting with STD in health facilities assessed and treated in an appropriate way (according to national standards).

Year	Area	Age	Rate	N=
1994	All		5.5	

Comments:

Sources: PPI 6&7 Survey. National AIDS Control Programme/WHO 1994

8 – United Republic of Tanzania

Health service indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. These efforts may range from reaching out to vulnerable communities through large-scale educational campaigns or interpersonal communication; provision of treatment for STIs; distribution of condoms and needles; creating and enabling environment to reduce risky behaviour; providing access to voluntary testing and counselling; home or institutional care for persons with symptomatic HIV infection; and preventing perinatal transmission and transmission through infected needles or blood in health care settings. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators – such as the percentage of a population with access to health care services; the percentage of women covered by antenatal care; or the percentage of immunized children – may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS – related issues.

☐ Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services – total:			
% of population with access to health services – urban:			
% of population with access to health services – rural:			
Contraceptive prevalence rate (%):	1990-1999	18	UNICEF/UNPOP
% of births attended by trained health personnel:	1990-1999	38	UNICEF
% of 1-yr-old children fully immunized – DPT:	1995-1998	74	UNICEF
% of 1-yr-old children fully immunized – Polio:	1995-1998	75	UNICEF
% of 1-yr-old children fully immunized – Measles:	1995-1998	72	UNICEF
Proportion of blood donations tested:			
% of ANC clinics where HIV testing is available:			
HIV/AIDS Hospital Occupancy Rate (Days):			

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programmes implement activities to increase both availability of and access to condoms. The two condom availability indicators below are intended to highlight areas of strength and weakness at the beginning and end of the distribution system so that programmatic resources can be directed appropriately to problem areas.

☐ Condom availability (central level)

Prevention Indicator 2: Availability of condoms in the country over the last 12 months (central level).

Year	Area	N	Rate
1997	All		2.8

Comments:

Sources: Population Service International, Family Planning Unit and National AIDS Control Programme

☐ Condom availability (peripheral level)

Prevention Indicator 3: Proportion of people who can acquire a condom (peripheral level).

Year	Area	N	Rate
1997	All		100.0

Comments:

Sources: Population Service International, Family Planning Unit and National AIDS Control Programme

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, intravenous drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of regular behavioural surveys in order to monitor trends in behaviours and target interventions.

☐ Knowledge of HIV- related preventive practices

Prevention Indicator 1: Proportion of people citing at least two acceptable ways of protection from HIV infection.

Year	Area	Age Group	Male	Female	All
1996	All	15-19	78.6	65.1	
1996	All	15-49	76.1	76.4	76.3
1996	All	20-24	76.6	81.0	80.1
1996	All	25-49	75.9	78.4	77.9

Comments:

Sources: :

☐ Reported non-regular sexual partnerships

Prevention Indicator 4: Proportion of sexually active people having at least one sex partner other than a regular partner in the last 12 months.

Year	Area	Age Group	Male	Female	All
1990	All	15-19	80.5	39.1	
1996	All	15-19	25.7	11.6	
1990	All	15-49	45.4	19.3	
1996	All	15-49	29.1	12.9	
1990	All	20-24	67.9	20.0	
1996	All	20-24	49.4	18.3	
1990	All	25-39	31.5	15.1	
1996	All	25-49	24.6	12.1	
1990	All	40-49	23.0	10.3	

Comments:

Sources:

☐ Reported condom use in risk sex (gen pop)

Prevention Indicator 5: Proportion of people reporting the use of a condom during the most recent intercourse of risk.

Year	Area	Age Group	Male	Female	All
1996	All	15-19	25.4	18.7	15.9
1996	All	20-24	40.3	21.9	29.1
1996	All	25-49	33.8	14.4	22.1

Comments:

Sources: Tanzania Demographic Health Surveys TDHS, 1996

10 – United Republic of Tanzania

Knowledge and behaviour

☐ Ever use of condom

Percentage of people who ever used a condom.

Year	Area	Age Group	Male	Female	All
1992	All	15-19		1.8	
1996	All	15-19	10.0	3.5	
1992	All	20-24		6.2	
1996	All	20-24	26.8	10.7	
1992	All	25-29		3.7	
1996	All	25-29	26.6	8.9	
1992	All	30-34		4.8	
1996	All	30-34	25.7	8.3	
1992	All	35-39		3.7	
1996	All	35-39	20.5	6.7	
1992	All	40-44		2.6	
1996	All	40-44	14.3	5.0	
1992	All	45-49		0.8	
1996	All	45-49	8.1	1.9	
1996	All	50-54	9.6		
1996	All	55-59	5.5		
1992	All	Total		3.6	
1996	All	Total		7.0	

Comments:

Sources: Demographic and Health Survey

☐ Median age at first sexual experience

Median age of people at which they first had sexual intercourse.

Year	Area	Age Group	Male	Female	All
1996	All	20-24	17.8	17.4	
1996	All	25-49	18.0	16.7	
1996	All	45-49	18.1	16.6	

Comments:

Sources: Tanzania Demographic Health Surveys TDHS, 1996

☐ Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child

Year	Area	Age Group	N	Rate
1996	All	15	288	1.3
1996	All	16	1039	10.3
1996	All	17	408	11.2
1996	All	18	343	39.2

Comments:

Sources: Tanzania Demographic Health Surveys TDHS, 1996

Sources

Data presented in this Epidemiological Fact Sheet come from several different sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations Agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

-
- Ali, A. K., O. J. Khatib, W. Osei, et al., 1992, Sentinel Surveillance for HIV Infection: Five Years Period, VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-12, Poster T.P.036.
- Hemed, Y., J. Minja, E. Naegele, et al., 1989, Seroepidemiological Study of Sexually Transmitted Agents (HIV, HBV, T. Pallidum) among Patients in the Mbeya Region (South Ta. ., IV Internat. Conf.: AIDS and Assoc. Cancers in Africa, Marseille, Oct. 18-20, Abstract 080.
- Haukenes, G., J. Shao, F. Mhalu, et al., 1992, The AIDS Epidemic in Tanzania: Rate of Spread of HIV in Blood Donors and Pregnant Women in Dar es Salaam, *Scandinavian Journal of Infectious Diseases*, vol. 24, pp. 701-706.
- Haji, S. H., A. K. Ali, A. A. Othman, et al., 1995, Prevalence of HIV Infection in Inmate Prisoners Pemba Island, Zanzibar, Tanzania, IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Abstract TuD128.
- Kipuyo, E. L., 1997, HIV-Trends and Behaviour Changes in Tanzania, Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Uganda", 10 - 13 February, Nairobi, Kenya.
- Laukamm-Josten, U., D. Ocheng, B. K. Mwizarubi, et al., 1992, HIV and Syphilis Seroprevalence and Risk Factors in Truckstops and Nearby Communities in Tanzania, VIII International Conference on AIDS, Amsterdam, 7/19-24, Poster PoC 4162.
- Lisekie, F., H. Grosskurth, A. Klokke, et al., 1992, Comparison of Sentinel and Cross-Sectional Study Data: Is Sentinel Surveillance a Useful Tool to Monitor the HIV/AIDS Epidemic?, VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster T.P.035.
- Lyamuya, E. F., E. Urassa, E. Mbena, et al., 1996, Stabilization of HIV-1 Seroprevalence in Pregnant Women in Dar es Salaam, Tanzania: Towards Reaching a Plateau Phase?, XI International Conference on AIDS, Vancouver, 7/7-14, Poster Tu.C.2463.
- Mhalu, F., U. Bredberg-Raden, E. Mbena, et al., 1987, Prevalence of HIV Infection in Healthy Subjects and Groups of Patients in Tanzania, *AIDS*, vol. 1, no. 4, pp. 217-221.
- Mhalu, F., M. Ngaiza, A. B. M. Swai, et al., 1991, Factors Associated with HIV Infection among a Sample of Bar and Restaurant Workers in Dar es Salaam, Tanzania, VII International Conference on AIDS, Florence, Italy, 6/16-21, Poster W.C.3087.
- Mwakagile, D. S. M., A. B. M. Swai, S. Mwambinga, et al., 1991, Socio-Epidemiological and Microbiological Aspects of Sexually Transmitted Diseases and HIV Infection in Dar es Salaam, Tanzania, VI International Conference on AIDS in Africa, Dakar, Senegal, 12/16-19, Abstract W.A.144.
- Matasha, E., J. Changalucha, H. Grosskurth, et al., 1992, Commercial Sexual Workers Intervention Programme: A Pilot Project in Northern Tanzania: Operational Data and International. ., VIII International Conference on AIDS, Amsterdam, 7/19-24, Poster PoD 5639.
- Mwakagile, D. S. M., A. B. M. Swai, K. J. Pallangyo, et al., 1992, Trend of Anogenital Warts among Patients Seen at a Referral Clinic for Sexually Transmitted Diseases in Dar es Salaam, Tanzania, VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster W.P.190.
- Mhalu, F., A. Swai, D. Mwakagile, et al., 1992, Surveillance and Control of HIV-1 Transmission among Female Bar workers in Dar es Salaam 1986-1991, VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster T.P.108.
- Mwakagile, D., A. Swai, J. Kitange, et al., 1993, Epidemiology of STDs in Dar es Salaam, Tanzania, VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Session M.O.P.058.
- Mwakagile, D., R. Mkuna, A. B. M. Swai, et al., 1993, Diagnosis of Sexually Transmitted Diseases (STDs) in Pregnant Women, VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Abstract M.O.P.027.
- Mwizarubi, B. K., K. Nyamuryekung'e, U. Laukamm-Josten, et al., 1995, Risk Factors for HIV Infection among Truck Drivers and Their Sexual Partners in Tanzania: Compilation of Cross-Sectional. ., IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Poster TuC579.
- Mwambingu, S. H., G. Rutahindurwa, F. Mhalu, et al., 1995, Development of STD Services in an Urban Clinic in Dar es Salaam, Tanzania, IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Poster MoB452.
- Msamanga, G. I., E. Urassa, D. Spiegelman, et al., 1996, Socioeconomic Status and Prevalence of HIV Infection among Pregnant Women in Dar es Salaam, Tanzania, XI International Conference on AIDS, Vancouver, 7/7-14, Poster Tu.C.2464.
- Mwakagile, D., A. B. M. Swai, E. Sandstrom, et al., 1996, High Frequency of Sexually Transmitted Diseases among Pregnant Women in Dar es Salaam, Tanzania: Need for Intervention, *East African Medical Journal*, vol 73, no. 10, pp. 675-679.
- Mwakagile, D., E. Mmari, C. Makwaya, et al., 1998, Unusual Characteristics of Adolescents and Youths Attending a New Special STD Clinic for Them in Dar es Salaam; . ., 12th World AIDS Conference, Geneva, 6/28 - 7/3, Abstract 22203.
- Nkya, W. M. M. M., W. P. Howlett, C. Assenga, et al., 1987, Seroepidemiology of HIV-1 Infection in the Kilimanjaro Region, *AIDS and Associated Cancers in Africa*, Naples, pp. 61-65.
- Nkya, W. M. M. M., W. P. Howlett, C. Assenga, et al., 1988, AIDS Situation in Northern Tanzania, III International Conference: AIDS and Associated Cancers in Africa, Sept. 14-16, Poster TP 9.
-

Websites: www.aids.africa.com

12 – United Republic of Tanzania

Annex: HIV Surveillance data by site

Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Pregnant women	Major Urban Areas	Dar es Salaam(1)		3.7		7.8	8.9	8.9	9.1	11	10.6	13.8	7.3				15.3
		Dar es Salaam(2)							10.3		16.1		12.2				14.1
		Dar es Salaam(3)									15.2		13.7				18.1
		Dar es Salaam(4)									15.3						
Pregnant women	Outside Major Urban Areas	Arusha				5.5					1						
		Bukoba						22.2	20	27.7	17.3						
		Dodoma				3											
		Iringa (Mafinga)							21	25							20.9
		Kagera (Bukoba)															7.0
		Kilimanjaro (Moshi)														19.9	
		Kilimanjaro (Umbe)							2.3	6.4	15.8		0	9.1	10		18.8
		Lindi					0.4										
		Mara (Musoma)							5.9	6.5	7.7						
		Mara (Nyasio)							9	7	8		7.2				
		Mbeya (Chimala)				4.2	6.3	8.8	9.5	8	10.8	16	10.5	17	16		
		Mbeya (Isoko)				2.9	2	2.4	6.6	18	8.5	8	10.3	7.5	7.3		
		Mbeya (Itete)				1.7	9.1	6.4	3.9	5.3	15.5	5.1	14.8	5.5	14.8		
		Mbeya (Kiwanjampaka)				7	10.6			17	22.3	19.5			22		
		Mbeya (Kyela)					21.2	14.6	17.5	30.4	27.2	27.5	32.5	23.1	44.4	24	
		Mbeya (Mbozi)											13.9	17	24		
		Mbeya (Meta)				10.3	16.9			25	13.7	16			32		
		Mbeya (Mwanbani)					12	8.5	12.9	8	10.7	13	17.5	16	14		
		Mbeya (Mwanjelwa)				11	7.3			11	23.3	19.6			36		
		Mbeya (rural)														12.3	
		Mbeya (urban)												18.7		17.3	
		Morogoro (urban)															18.4
		Morogoro (Turiani)															9.5
		Mtwara (Nanguruwe)							4.4				0				
		Mwanza				6											
		Mwanza (Mkula)							3.7	4.6	5.4		5.4				
		Mwanza (rural)							3.7	3.9	5.5	5.5	5.4				
		Mwanza (urban, 1)				8	11.7	12.2	15.3	10	12.3	11.7	20.3				
		Mwanza (urban, 2)						12.3	11.2				8.9				
		Mwanza (urban, 3)											8.7				
		Pemba Island						0.3	0	0.6	0.8	0.3					
		Rukwa (Namanye)								11.3	8.3			11.2	11.2		
		Rukwa (Sumbawange)							12	23.2			22.2		21		
		Ruvuma (Madaba)							0.6	12.9							
		Ruvuma (Namtumbo)							3.5	6.7	3.2	5.6			4		
		Ruvuma (Songea)							9.7	16.1	15.7	14.2			9.8		
		Singida				3.6											
		Tanga				7											
		Zanzibar				0.3	0.5	0.6	0.6	0.7	0.5	2.3					
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Sex workers	Major Urban Areas	Dar es Salaam		29		42.1	43.7	45	42.9		49.5						
Sex workers	Outside Major Urban Areas	Arusha				18.6											
		Dodoma				19.5											
		Kilimanjaro				14.8											
		Morogoro									60.9						
		Moshi				31.9											
		Mwanza town							39.8								
		Singida				7.5											
		Tanga				25.7											
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Injecting drug users	Major Urban Areas																
Injecting drug users	Outside Major Urban Areas																

12 continued – United Republic of Tanzania

Annex: HIV Surveillance data by site

Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI patients	Major Urban Areas	Dar es Salaam										24			13.2		
		Dar es Salaam (Males, 1)		9.3	13.4	21.8	28.3	24.5	16.1								
		Dar es Salaam (Males, 2)		13.2		27.8	39.7		18.5								
		Dar es Salaam (Males, 3)							14.4								
		Dar es Salaam (Females, 1)		12.2	28.1				21.3								
		Dar es Salaam (Females, 2)		27.8					40.4								
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI Patients	Outside Major Urban Areas	Kagondo (Males)			20												
		Mbeya				22.5		27.5	32	34.3							
		Mbeya (Chimala)													18		
		Mbeya (Females)				23.6											
		Mbeya (Isoko)													22		
		Mbeya (Itete)													27		
		Mbeya (Kiwanjampaka)													38		
		Mbeya (Kyela)													25		
		Mbeya (Males)					21.4										
		Mbeya (Mbozi)													43		
		Mbeya (Mwanbani)													36		
		Mbeya (Referral Hospital)													17		
		Mbeya (Ruanda municipal)													36		
		Zanzibar								5.2	28.2						
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Blood Donors	National																
Blood Donors	Major Urban Areas																
Blood Donors	Outside Major Urban Areas																